

Name:

Date:

Degree of water saturation

Description of the sample, material:

Specific density

$$\rho_s = \frac{m_s}{V_s} \quad [\text{Mg/m}^3]$$

$m_1 = \dots\dots\dots \text{ g } \dots\dots\dots \text{ g } \dots\dots\dots \text{ g}$

$m_2 = \dots\dots\dots \text{ g } \dots\dots\dots \text{ g } \dots\dots\dots \text{ g}$

$m_3 = \dots\dots\dots \text{ g } \dots\dots\dots \text{ g } \dots\dots\dots \text{ g}$

$$\rho_s = \frac{m_2 - m_1}{m_4 - m_1 + m_2 - m_3} \cdot \rho_k \quad [\text{Mg/m}^3]$$

$m_4 = \dots\dots\dots \text{ g } \dots\dots\dots \text{ g } \dots\dots\dots \text{ g}$

T = °C → $\rho_k = \dots\dots\dots \text{ Mg/m}^3$

$\rho_s = \dots\dots\dots \text{ Mg/m}^3$

m_1 – pycnometer weight

m_2 – weight of pycnometer and dried sample

m_3 – weight of pycnometer and sample with liquid

m_4 – weight of pycnometer with liquid

ρ_k – density of liquid (temperature dependent)

$\rho_s = \dots\dots\dots \text{ Mg/m}^3$

Water content

$$w = \frac{m_w}{m_s} \cdot 100 \quad [\%]$$

$m = \dots\dots\dots \text{ g } \dots\dots\dots \text{ g } \dots\dots\dots \text{ g}$

$m_s = \dots\dots\dots \text{ g } \dots\dots\dots \text{ g } \dots\dots\dots \text{ g}$

$m_w = m - m_s \quad [\text{g}]$

$m_w = \dots\dots\dots \text{ g } \dots\dots\dots \text{ g } \dots\dots\dots \text{ g}$

$w = \dots\dots\dots \% \dots\dots\dots \% \dots\dots\dots \%$

m – total weight of the sample

m_w – weight of the water in the sample

m_s – weight of the dried sample

$w = \dots\dots\dots \%$

Bulk (total) density REGULARLY SHAPED SAMPLE

$$\rho = \frac{m}{V} \quad [\text{Mg/m}^3]$$

$m = \dots\dots\dots \text{ g}$

$a \text{ (d)} = \dots\dots\dots; \dots\dots\dots; \dots\dots\dots; \text{ average: } \dots\dots\dots \text{ mm}$

$b = \dots\dots\dots; \dots\dots\dots; \dots\dots\dots; \text{ average: } \dots\dots\dots \text{ mm}$

$h = \dots\dots\dots; \dots\dots\dots; \dots\dots\dots; \text{ average: } \dots\dots\dots \text{ mm}$

$V = \dots\dots\dots \text{ mm}^3 = \dots\dots\dots \text{ m}^3$

m – weight of the sample (natural moisture)

$a \text{ (d)}$ – width of block (diameter)

b – length of block

h – height of block

$\rho = \dots\dots\dots \text{ Mg/m}^3$

Bulk (total) density IRREGULARLY SHAPED SAMPLE

$$\rho = \frac{m}{V} \quad [\text{Mg/m}^3]$$

m = g g g

m₁ = g g g

$$V = \frac{m_1 - m_2}{\rho_w} - \frac{m_1 - m}{\rho_p} \quad [\text{m}^3]$$

m₂ = g g g

$\rho_w = 1 \text{ Mg/m}^3$

$\rho_p = 0.9 \text{ Mg/m}^3$

V = m³ m³ m³

$\rho = \text{..... Mg/m}^3$

m – weight of the sample (natural moisture)

m₁ – weight of the sample with paraffin

m₂ – weight of the sample with paraffin under water

ρ_w – density of water (temperature dependent)

ρ_p – density of paraffin

$\rho = \text{..... Mg/m}^3$

Degree of water saturation

$$S_r = \frac{V_w}{V_p} \cdot 100 \quad [\%]$$

$$S_r = \frac{w \cdot \rho \cdot \rho_s / \rho_w}{\rho_s \cdot (w + 1) - \rho} \quad [\%]$$

$S_r = \text{..... } \%$